

DR. SHARON DEMORROW (Orcid ID : 0000-0002-0511-9778)

DR. CHRISTOPHER F ROSE (Orcid ID : 0000-0001-9854-6834)

Article type : Reviews

Editor : Luca Valenti

2021 ISHEN Guidelines on Animal Models of Hepatic Encephalopathy

Authors:

¹DeMorrow S*,

²Cudalbu C,

³Davies N,

⁴Jayakumar AR,

⁵Rose CF

Affiliations:

¹ Division of Pharmacology and Toxicology, College of Pharmacy, The University of Texas at Austin, Texas, USA; Department of Internal Medicine, Dell Medical School, The University of Texas at Austin, Texas, USA; Research division, Central Texas Veterans Healthcare System, Temple Texas USA.

² Center for Biomedical Imaging, Ecole Polytechnique Fédérale de Lausanne, Lausanne, Switzerland

³ Institute for Liver and Digestive Health, University College London, Royal Free Campus, London, United Kingdom

⁴ General Medical Research, Neuropathology Section, R&D Service and South Florida VA Foundation for Research and Education Inc; Obstetrics, Gynecology and Reproductive Sciences, University of Miami School of Medicine, Miami FL, USA.

⁵ Hepato-Neuro Laboratory, CRCHUM, Université de Montréal, Montreal, Canada

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/LIV.14911](https://doi.org/10.1111/LIV.14911)

This article is protected by copyright. All rights reserved

***Correspondance:** Sharon DeMorrow, PhD, Sharon.demorrow@austin.utexas.edu; tel: +1-512-495-5779

Word count: ~5000

Total number of Figures and Tables: 4 figures, 2 tables

List of Abbreviations:

ISHEN, International Society of Hepatic Encephalopathy and Nitrogen Metabolism; HE, hepatic encephalopathy; ALF, acute liver failure; CLD, chronic liver disease; AOM, Azoxymethane; APAP, Acetaminophen; TAA, Thioacetamide; CCl₄, Carbon tetrachloride; D-GAL, Galactosamine; BDL, bile-duct ligation; PCA, portacaval anastomosis; NADP⁺/NADPH, reduced and oxidized forms of nicotinamide adenine dinucleotide phosphate; AST, aspartate transaminase; ALT, alanine transaminase; ICP, intracranial pressure; TIPS, transjugular intrahepatic portosystemic shunt; LPS, Lipopolysaccharide; OP, Ornithine-phenylacetate; BBB, blood brain barrier.

Conflict of Interest Statement:

SD: nothing to disclose

CC: nothing to disclose

ND: nothing to disclose

ARJ: nothing to disclose

CFR: Christopher F. Rose has research collaborations with Mallinckrodt Pharma and Neuractas and is an advisor for Axcella, Horizon Therapeutics, Lupin Pharma, Morphocell Technologies, and Neuractas.

Statement of Financial support:

Financial support was provided by the Swiss National Science Foundation (project no 310030_173222)

Acknowledgements: Parts of this review were prepared with resources and the use of facilities at the Central Texas Veterans Health Care System, Temple, Texas. The content is

the responsibility of the author(s) alone and does not necessarily reflect the views or policies of the Department of Veterans Affairs or the United States Government.

Parts of this work was also made possible thanks to the Center for Biomedical Imaging (CIBM), founded and supported by Lausanne University Hospital (CHUV), University of Lausanne (UNIL), Ecole Polytechnique Federale de Lausanne (EPFL), University of Geneva (UNIGE) and Geneva University Hospitals (HUG)

Abstract:

This working group of the International Society of Hepatic Encephalopathy and Nitrogen Metabolism (ISHEN) was commissioned to summarize and update current efforts in the development and characterization of animal models of hepatic encephalopathy (HE). As defined in humans, HE in animal models is based on the underlying degree and severity of liver pathology. Although hyperammonemia remains the key focus in the pathogenesis of HE, other factors associated with HE have been identified, together with recommended animal models, to help explore the pathogenesis and pathophysiological mechanisms of HE. While numerous methods to induce liver failure and disease exist, less have been characterized with neurological and neurobehavioral impairments. Moreover, there still remains a paucity of adequate animal models of Type C HE induced by alcohol, viruses and non-alcoholic fatty liver disease; the most common etiologies of chronic liver disease.

Key words: hepatic encephalopathy, acute liver failure, chronic liver disease, liver toxins, hyperammonemia, neurological deficits, behavior, bile duct ligation, portocaval anastomosis.

Key points

- Valid animal models of liver failure/disease which manifest features/elements of HE currently exist.
- A suitable animal model of HE must have; (i) some degree of liver injury, failure and/or vascular impairment, (ii) hyperammonemia and (iii) some degree of neurological, behavioral or motor impairment.
- New animal models are warranted with a particular emphasis on the common etiologies associated with chronic liver disease (alcohol, hepatic viral infections and fatty liver disease) and their impact on the development of HE.
- An appropriate animal model should be selected once the interested feature(s) of HE have been identified.

Introduction:

Hepatic encephalopathy (HE) is a debilitating neurological complication of liver disease/failure characterized by cognitive, psychiatric and motor disturbances. Due to existing animal models of HE, many new insights into the pathogenesis and pathophysiology of this disorder have been realized. In addition, many animal models have been used for pre-clinical studies testing new therapeutic strategies for HE. Three types of HE have been defined based on degree and severity of liver failure, disease or impairment (including degree of portal-systemic shunting); (1) Type A HE: associated with acute liver failure (ALF) resulting from the rapid onset of hepatocyte necrosis and severe inflammation without pre-existing liver impairment; (2) Type B HE: associated with portal-systemic shunting in the absence of liver disease or failure; (3) Type C HE: associated with chronic liver disease (CLD; severe fibrosis or cirrhosis). A common feature of the three types of HE is that the overall capacity of the liver to remove ammonia is reduced and hyperammonemia ensues.

What defines a good model of HE?

The fundamental basis for an animal model of HE is the presence of liver injury, failure and/or impairment that consequently leads to hyperammonemia. In addition, an animal model of HE should also exhibit some degree of neurological, behavioral or motor impairments. Other features believed to be implicated in HE include systemic inflammation¹ and oxidative stress² as well as neuroinflammation³⁻⁵. However, their roles in the pathogenesis of HE in the absence of hyperammonemia as well whether they are contributory factors or consequential effects remains undetermined. The cardinal features of a Type A HE animal model is progression to overt neurological symptoms such as loss of corneal reflex (coma) and loss of righting reflex (pre-coma) within hours or days upon induction of ALF. Moreover, intracranial hypertension is another HE landmark associated with ALF which depicts cerebral edema, which is commonly observed in most animal models of Type A HE. In addition, various scoring scales have been developed for HE in ALF based on observed behavioral changes, impairment in reflexes and/or the presence of ataxia^{6,7}. A commonly used methodology involves assigning the animals to a particular stage of HE

based on their observed behaviors, with the data presented as the average stage of HE per experimental cohort at any time point⁶. Recently, an improved categorical neurological scoring system was developed whereby 5 individual reflexes and presence of ataxia were each assessed and assigned a score from 0-2 (2 = reflexes intact, no ataxia; 1 = reflexes delayed, minor ataxia, 0 = reflexes absent, significant ataxia). The neurological score is the sum of scores for each reflex at any time point⁷ (Table 1). Type B HE, as well as Type C HE, both chronic models of HE, do not develop evident or overt neurological symptoms (including obvious behavioral changes, reflex impairments). As a result, HE is evaluated through different neurobehavioral tests⁸ such as Open Field, Morris Water Maze and Y-Maze tests which assess exploratory behavior, locomotor activity, anxiety and memory (including spatial and working), respectively (Table 2). Furthermore, while an increase in intracranial pressure (ICP) is rare in Type C HE, brain edema is still present. However, whether low-grade edema or age-related brain atrophy prevents an increase in ICP in patients with CLD still remains unresolved⁹.

Various degrees of liver failure/injury and hepatic vascular impairment can be induced by i) numerous liver toxins (including azoxymethane (AOM), acetaminophen (APAP), thioacetamide (TAA), carbon tetrachloride (CCl₄) and galactosamine (D-GAL)), with the dose (+/- barbiturate pre-treatment to enhance the sensitivity to the toxin), and length of treatment dictating the type of HE model: Type A vs. Type C and ii) surgical interventions (bile-duct ligation (BDL), portacaval anastomosis (PCA) and PCA + hepatic artery ligation (HAL)) (Figure 1). Whereas the major etiologies of CLD in humans include alcohol, viral hepatitis and non-alcoholic fatty liver disease (NAFLD), for which appropriate animal models of severe fibrosis/cirrhosis currently do not exist. This is primarily due to the resistance of rodents to such etiological factors which do not develop severe fibrosis/cirrhosis. Consequently, this remains a major obstacle in the field which hinders the understanding of the independent role of each of the etiological factors on brain function.

Sampling and measuring ammonia:

Since elevated blood ammonia levels are a cardinal feature of HE, measuring ammonia is imperative and various methods to quantify ammonia currently exist. The most commonly used method to measure ammonia involves an enzymatic kinetic assay in which ammonia reacts with α -ketoglutarate and nicotinamide adenine dinucleotide phosphate (reduced form NADPH) to form glutamate and NADP⁺. The amount of ammonia is related to the amount of NADPH oxidized, which is measured photometrically. Regardless of the method, the measurement of ammonia is extremely sensitive, with the stability of the molecule being an important issue which needs to be carefully controlled. As a result, anomalous plasma ammonia concentrations can be due to pre-analytical events such as delayed sample processing, temperature, type of test matrices, hemolysis and storage time in freezer^{10,11}. Accordingly, it is important to place the sample on ice and immediately remove the plasma from cells to reduce the reaction rate of ammonia metabolism. The delay between sampling and analysis should be kept within 2hrs¹². Heparin, EDTA and oxalate anticoagulated plasma have been tested with an EDTA-anticoagulated matrix found to be superior¹³. Frozen samples have been shown to affect ammonia levels when extensively kept frozen as well as multiple freezing and thawing cycles having a profound effect. Measuring brain tissue ammonia is further challenging due to the multiple pre-analytical steps including extraction of the brain and post-mortem metabolic changes. Aside from using liquid nitrogen, brains can be fixed for analysis by using focused beam microwave irradiation. This is a technique which causes permanent inactivation of enzymes in less than 1s, thereby minimizing enzyme-dependent post-mortem metabolic changes¹⁴. Nessler's staining is a histological method to identify high levels of ammonia (corresponding plasma levels $\geq 150\mu\text{M}$) which has been demonstrated in liver¹⁵. However, this technique has not been validated in brain. Alternatively, cerebrospinal fluid and microdialysate measurements of ammonia have been performed which are believed to better reflect brain ammonia concentrations¹⁶.

Current animal models of HE:

There are a number of advantages and disadvantages in using small animals vs. large animals which may explain the choice of a particular animal model (Figure 2). Primarily

mice, rats and pigs have been used in recent years for studies in HE and therefore we will focus on these species. Type A HE have been developed to induce ALF and subsequent overt, progressive neurological decline. In a majority of these models care must be taken to control associated features such as body temperature, blood glucose and potassium levels to prevent these confounders from interfering with outcomes and interpretation of data. Models of Type B and C HE mimic features of covert/minimal HE where overt symptoms do not develop and are not observed.

Type A HE:

Liver toxin models:

Acetaminophen:

Rodents: Also known as paracetamol, APAP is commonly used clinically as an antipyretic and analgesic. APAP is metabolized by p450s leading to N-acetyl-p-benzoquinoneimine (NAPQI) when intracellular glutathione levels are saturated. Therefore, high doses of APAP lead to centrilobular hepatocyte necrosis. While the APAP-treated experimental model of Type A HE was not well established prior to the 2009 ISHEN guidelines¹⁷, there is substantial use of APAP to induce Type A HE both in rats and mice. APAP is injected at various dosages (from 300 mg/kg – 5 g/kg, ip) with mice being more susceptible than rats¹⁸. APAP-induced ALF in mice leads to brain edema and coma together with increased blood ammonia levels¹⁹. Additionally, activation of intracellular signaling mechanisms, including elevated levels of cytokines, activation of brain transcription factors (increased levels of NF- κ B, Nrf2 etc.), induction of oxidative stress, glial fibrillary acidic protein expression, as well as autophagy-related events have been observed in the brains of these animals¹⁹⁻²⁷.

Pigs: Large animal models of ALF have been a long-standing topic of interest for testing of 'human scale' therapeutic devices. However, there was considerable difficulty in creating a stable, reproducible model that reflected the development of human toxicity²⁸. In 2011 work by Thiel et al. demonstrated that a reproducible model of ALF can be achieved through careful monitoring of blood APAP levels within a narrow window (300-450 mg/dL)²⁹, which was further refined by Lee *et al.* to show that effective studies could be conducted in

relatively few animals³⁰. This APAP-induced ALF model demonstrates core features of disease as a consequence of progressive hepatic necrosis and the consequent systemic inflammatory response. Hyperammonemia develops progressively once liver failure is established to levels in excess of 300 μ M at exitus, concomitant with the occurrence of increased intracranial pressure (ICP) elevation (>30 mmHg)²⁹. The model is conducted in a terminally anesthetized animal with continuous intensive care support, preventing testing of any behavioral or cognitive impairments. However, the availability of samples and the ability to use invasive tools to investigate specific mechanisms has allowed a better understanding of the pathophysiology of the condition³¹⁻³³.

Azoxymethane:

Rodents: AOM is an active metabolite of the cycad palm nut (found primarily on the island of Guam) which is hepatotoxic. The AOM model of ALF has been used to characterize the molecular changes associated with the development of HE in mice. It involves a single intraperitoneal (ip) injection of AOM at doses ranging from 50 mg/kg to 100 mg/kg^{6,7,34,35}. Characterized by massive hepatocyte cell death, metabolism of AOM by cytochrome p450s generates toxic metabolites in the liver resulting in the formation of DNA adducts³⁶. Associated with acute severe liver injury is a progressive loss of reflexes, acquisition of cognitive dysfunction, and ultimately hepatic coma^{6,7,34} within 24hr to 48hr. It is associated with hyperammonemia^{6,34} cerebral edema^{6,34,37}, presence of neuromuscular deficits (manifested by ataxia), microglia activation and the upregulation of neuroinflammatory signals^{6,7,34}. Additionally, studies demonstrating AOM induces ALF and HE in rats are lacking.

Galactosamine:

Rodents: Type A HE animal models using D-GAL have been reported since 1970. D-GAL is an amino sugar, which when metabolized in the liver consequently disturbs hepatocyte RNA metabolism, resulting in hepatocyte necrosis. The D-GAL model of Type A HE in rats replicates characteristic features of HE including the characteristic stages of HE, altered liver

enzymes and histopathological changes similar to that observed in humans with Type A HE^{38,39}. D-GAL given at a dose of up to 1.5g/kg (single dose, ip) leads to severe hepatic necrosis (diffuse degenerative changes with focal coagulative necrosis and the appearance of eosinophilic bodies), inflammatory infiltration of periportal areas and serum levels of liver enzymes are greatly increased while prothrombin is reduced⁴⁰. There is a vast array of studies demonstrating diverse effects with different doses of D-GAL with 2.5g/kg (ip) leading to progressive HE and death within 48hrs, accompanied with an increase in ICP⁴¹. D-GAL induced HE studies in mice are few and therefore lacking. Furthermore, the limitations of this model include a lack of reproducibility since animals die unexpectedly and even brain edema, increased ICP and coma are not consistently observed.

Pigs: An alternative large animal of Type A model that has attracted attention in recent years is that of a bolus injection of D-GAL in pigs (0.75g/kg iv)⁴². This model has demonstrated the core symptoms of Type A HE within 48 hours, with blood biochemistry meeting the criteria for ALF, elevated plasma ammonia levels and raised ICP (>25mmHg) in combination with histological evidence of ALF obtained from sequential biopsies⁴². Animals are supported throughout the study with dextrose and crystalloid solutions as required and without further intervention, survival is expected to be between 72-90 hours. There are several advantages to using this model including the ability to study awake animals and observe the development of the stages of HE progressing to pre-coma and eventually coma, at which time point the animals are euthanized. However, it should be noted that a high degree of variability has also been found in this model⁴³ and there have been questions raised as to its clinical relevance²⁸.

Thioacetamide:

Rodents: TAA's toxicity is induced by the production of thioacetamide-s-oxide (a reactive oxygen species) through flavin adenine dinucleotide monooxygenase⁴⁴. The TAA-treated rat model of Type A HE has been well-established relative to the clinical status, liver function, and brain edema development⁴⁵⁻⁵². In this model, rats are given TAA in doses ranging from 300 to 500 mg/kg by ip injection on consecutive days for 2-3 days. While not as well

characterized in mice, TAA-induced hepatotoxicity has shown to induce ALF and subsequent HE following the administration of TAA (100-300 mg/kg, ip) for 3 consecutive days⁵³⁻⁵⁶. TAA treatment in both species results in severe hepatocellular and bridging necrosis without cholestasis^{47,49}. Furthermore, increased blood and brain ammonia have been observed⁴⁹. Associated with the decline in neurological function and brain edema⁵⁴ are enlarged, vacuolated nuclei and pale/expanded cytoplasm in the astrocytes of the cerebral cortex⁵⁴ with the neuropil highly vacuolated (spongiotic), especially around blood vessels and neurons⁵⁴.

Surgical Models:

Hepatic devascularization:

Rodents: Hepatic devascularization in rats is an established model of Type A HE which has been commonly used since 2010⁵⁷⁻⁶⁰. Investigators have typically studied models of ALF by employing anastomosis of the portal vein to the vena cava and ligating the hepatic artery, seizing all blood flow to the liver. This model manifests a reproducible progression of Type A HE. A consistent and progressive increase in blood and brain ammonia as well as astrocyte swelling and brain edema have been characterized in this model⁶¹. Altered gene expression and CNS inflammation have also been identified^{62,63}. Hepatic devascularization in smaller rodents is technically challenging and therefore is yet to be developed or characterized in mice.

Pigs: There is no standardization in the literature on the use of hepatic devascularization models in large animals. Studies have reported the use of additional hepatic artery occlusion resulting in total devascularization with⁶⁴ or without reversal of the procedure⁶⁵. Other investigators have combined portal shunting with varying degrees of hepatectomy⁶⁶⁻⁶⁹ to create Type A models. As such, these models display a degree of heterogeneity in their findings and a spectrum of severities⁷⁰ from to intensively monitored terminal models limited to several hours post procedure to those with a recover phase⁷¹. Porcine models of hepatic devascularization demonstrate central features of Type A HE, including cerebral edema, intracranial hypertension, hyperammonemia and blood brain barrier (BBB) breakdown⁷²⁻⁷⁴,

along with hyperlactatemia and significant hemodynamic perturbation^{75,76}. Due to the fact that the majority of studies have been conducted in anesthetized models, there is scarce information on cognitive impairment in these models other than the reported changes in ICP⁷³, and the lack of a standardized approach to experimental design makes direct comparison difficult.

Type B HE:

Portacaval anastomosis:

Portal-systemic shunting (when the normal flow from the portal vein is diverted, either partially or completely, to the systemic circulation, thus bypassing the liver) leads a decrease in hepatic ammonia extraction and consequently the development of hyperammonemia, even in the absence of liver disease. Congenital shunts⁷⁷ occur in humans but are commonly observed in dogs which present with hyperammonemia and psychomotor dysfunction⁷⁸ and are often admitted to veterinarian clinics with various symptoms, including behavioral changes. Congenital shunts are believed to be strain specific as has been reported in C57BL/6J mice⁷⁷. In rats, an end-to-side portacaval anastomosis is a surgical procedure which leads to a rise in blood ammonia levels and neurological impairment⁷⁹. The diversion of blood during portal systemic shunting consequently leads to liver atrophy⁸⁰⁻⁸². These surgical procedures are also achievable in larger animals such as pigs⁸³ whereas it is very difficult to realize this surgery in mice. Alternatively, graded portal-vein stenosis which leads to spontaneous portal-systemic shunts and various degrees of portal-systemic shunting, is a much easier model to develop but with greater variability and therefore less reproducible⁸⁴. In CLD, an increase in hepatic resistance can lead to acquired liver shunts where the degree of shunting increases the risk of developing severe HE⁸⁵. Portal-systemic shunting without liver impairment is evidently not a model of liver disease, nor is it a model of transjugular intrahepatic portosystemic shunt (TIPS). Ideally, performing a portal-systemic shunt in an animal model of CLD would be a valuable model for studying post-TIPS HE.

Type C HE:

Liver toxin models:

Carbon Tetrachloride:

Rodents: CCl₄ is mainly metabolized by centrilobular hepatocytes producing the toxic metabolite trichloromethyl (CCl₃⁻) via cytochrome p-450s, causing centrilobular liver damage. CCl₄ administration in rodents vary in terms of route of administration (injections (ip, sc), gavage), dosage, frequency of dosing and duration. Long term treatment of rats or mice with CCl₄ leads to repeated insults to the liver causing hepatocyte damage, ductular reaction and myofibroblast activation, hepatic stellate cell activation, imbalance between extra cellular matrix production and degradation and development of progressive liver fibrosis^{86,87}. In mice, doses used range from 0.5 mL/kg to 1 mL/kg, administered either via oral gavage or ip injection, twice per week for up to 16 weeks⁸⁸⁻⁹¹. These treatments result in hyperammonemia^{89,91}, hyperpermeability of the BBB⁹¹, increased neuroinflammatory signals⁸⁸⁻⁹⁰, microglia/glia cell activation^{88,89}, and increased GABA signaling⁸⁸. In rats, CCl₄ administration (0.1 - 0.2 mL/kg twice per week; ip for up to 5 months)^{87,92,93} results in hyperammonemia^{87,92}, impaired memory acquisition as determined using the Morris Water Maze^{87,93} and a decrease in hippocampal neurogenesis⁸⁷.

Thioacetamide:

Rodents: Similar to CCl₄, TAA primarily causes centrilobular hepatocyte damage⁴⁴. A TAA-treated rat model of Type C HE has recently been established relative to the clinical status, liver function, and behavioral and cognitive deficits^{94,95}. In this model, rats were given TAA (100 mg/kg, ip) for 10 consecutive days. Hepatic damage (ballooning degeneration, hydropic changes and the presence of eosinophilic bodies, affecting approximately 60-70% of the liver parenchyma) as well as increased serum ALT and AST levels were observed in this model. The presence of Alzheimer type II astrocytosis (predominantly in the cerebral cortex), behavioral abnormalities associated with cognitive dysfunction are observed, including drowsiness, decreased wakefulness, impaired attentiveness, decreased grooming and exploratory behavior⁹⁶ together with increased blood and brain ammonia^{94,95}. In

addition, many studies have administered 12 weeks of TAA by its addition to the drinking water⁹⁷⁻¹⁰². TAA-induced type C HE models in mice are lacking.

Surgical models:

Bile-Duct Ligation:

Rodents: The BDL rat model is the most widely used model of type C HE and has been shown to be a reproducible model of biliary CLD, simulating metabolic aspects of cirrhosis. In this model the common bile duct is ligated (2 or 3 points) and then resected to avoid reversibility. BDL rats survive to 6 to 8 weeks post ligation, develop liver failure, jaundice, portal hypertension, portal-systemic shunting, bacterial translocation and immune system dysfunction^{17,103-105}. The signs of CLD in these rats include increased levels of plasma bilirubin, liver enzymes (AST and ALT) and histological changes in the liver architecture (i.e. bile duct epithelial cell proliferation, disturbed cytoarchitecture, formation of septae between portal areas and a noticeable increase in collagen fibers)¹⁰⁶. BDL rats also develop hyperammonemia and show motor and learning deficits¹⁰⁵⁻¹⁰⁸. *In vivo* neuro-imaging studies performed on BDL rats have reproduced the changes observed in humans (i.e. increased brain glutamine and decrease in brain osmolytes as an osmotic response and other more subtle changes including a decrease in antioxidants and creatine)^{105,107,109-111}. The changes in water content (i.e. brain edema) measured in BDL rats are subtle^{107,111} (details on brain water measurements and brain edema can be found in the following reviews^{8,9,112}). Systemic oxidative stress, as a result of primary liver injury, combined with hyperammonemia was shown to stimulate an increase in brain water content in BDL rats⁷⁹. Inflammation (systemic and central) has not been thoroughly characterized in these rats and results to date remain controversial¹¹³⁻¹¹⁵.

In contrast, BDL mice have been primarily used to study cholestatic liver injury and liver inflammation. The survival rate of BDL mice is much less than BDL rats, with studies lasting between 5 days post ligation till 2 weeks^{116,117}. It is possible to prolong survival in BDL mice, but this requires intensive care (i.e., cage warming, wet food placed on the cage floor, and potentially dextrose supplementation subcutaneously)¹¹⁸. The BDL mice are characterized

by decreased social investigative behavior and increased immobility together with increased serum bilirubin, changes in liver enzymes (ALT) and the presence of portal based inflammatory cells in stained liver sections¹¹⁷. Beyond these observations, no features of HE have been observed, likely due to the short survival time of the mice.

Models of overt HE in chronic liver disease:

In order to study the precipitating factors of overt HE and hence the pathogenic factors involved in the acute neurological impairment in CLD, insults are used to induce overt HE in animals. Ammonia, a driver of a number of precipitating factors, has been administered to rats to induce an overt episode of HE. Exacerbating hyperammonemia through diet in BDL rats has also elucidated overt symptoms of HE¹¹³. The use of PCA to shunt blood away from the liver to establish hepatic insufficiency prior to the administration of a secondary insult has similarly been extensively studied^{79,119}. Whereas early studies sought to investigate impaired hepatic clearance of bacteria¹²⁰ post PCA, more recent studies have investigated episodic HE induced by repeated administration of endotoxin or ammonia which demonstrated neuronal loss and cerebral inflammation^{121,122}. Other studies in this model have explored the effects of simulated esophageal bleeding¹²³ as a source of hyperammonemia, and the effects of superadded ammonia on other body systems^{124,125}. Lipopolysaccharide (LPS) injected into BDL rats is used to mimic severe inflammation and infection; a precipitating factor of overt HE. This model is the creation of a single inflammatory event on a background of established liver injury, after which the animals are euthanized for sample collection. The BDL with a secondary insult of bacterial endotoxin (BDL + LPS) was originally described in 1999 by Harry et al.¹²⁶ in which the heightened response to the inflammatory insult was observed. Subsequently this BDL + LPS model was shown to demonstrate the key features of HE with evidence of cerebral inflammation¹¹⁴. However, the key point was that there was significant worsening of most measured parameters following the administration of endotoxin in a relatively short time frame (3 hours), indicating the possibility of a priming effect during CLD development leading to a hyper-responsiveness to the secondary insult.

Ammonia supplementation to induce hyperammonemia:

In order to study the effect of neurotoxicity from elevated blood ammonia levels without the impact of liver disease, hyperammonemia is induced in naïve rats through an ammonia supplemented standard diet (ammonia acetate; 20% w/w)¹²⁷ to study the effects of hyperammonemia on brain function and metabolism in absence of liver dysfunction. Hyperammonemia is induced within 10 days and studies have demonstrated rats can tolerate this ammonia-supplemented diet for up to 100 days. Ammonium acetate is much more effective in the diet as oppose to in the drinking water. Also, other ammonia salts, such as ammonium carbonate are not well tolerated. In these diet-induced models, pair-fed animals are highly recommended as controls. Cognitive impairment has been demonstrated after 7 days of ammonia-supplemented diet¹¹³. This is a simple, reproducible animal model of chronic hyperammonemia.

Preclinical models for HE treatments:

Preclinical models of HE are imperative for testing novel treatments for HE. Even though treating the liver disease itself may resolve HE, it is very difficult to improve the liver at end-stage liver disease. Therefore, novel therapeutic strategies for HE primarily target a pathogenic factor which is precipitating the observed neurological impairment. In the CLD setting, the BDL rat is an excellent model to assess the efficacy of ammonia-lowering strategies. AST-120 carbon microspheres, when administered by gavage, were shown to lower blood ammonia in BDL rats, was associated with a reduction in brain edema and HE symptomatic behavior¹⁰⁴. Ornithine-phenylacetate (OP) and liposome-supported peritoneal dialysis have also shown to be beneficial in lowering blood ammonia levels in BDL rats^{128,129}. The probiotic VIVOMIXX has been recently shown to attenuate hyperammonemia and improve both the performance in behavioral tests and the neurometabolic profile of BDL treated rats¹³⁰. OP has also been shown to lower blood ammonia in rats with portal-caval anastomosis¹²³. Interestingly, GABA-A receptor antagonists have shown to be beneficial in PCA rats as well as rats administered ammonia in the diet¹³¹. In ALF, L-ornithine-L-

aspartate, minocycline, N-acetylcysteine and hypothermia have all shown to be beneficial in liver devascularized rats^{60,132-134}. Similarly, OP was beneficial in lowering ammonia and ICP in liver devascularized pigs¹⁶. In the AOM mouse model, strategies to inhibit cerebral inflammation, such as targeting proinflammatory cytokine signaling, or inducing mild hypothermia were shown to attenuate parameters of HE^{3-5,135-137}. Furthermore, emerging evidence suggesting that aberrant bile acid signaling in the brain may contribute to the pathogenesis of HE in the AOM mouse model and that inhibition of bile acid receptors such as farnesoid X receptor or sphingosine-1-phosphate receptor 2 was recently shown to be beneficial¹³⁸⁻¹⁴⁰.

Concerns when using animal models of HE

Several limitations exist in using toxin models of both Type A and C HE. Firstly, there is controversy surrounding the reported off-target effects of the toxins that may confound the interpretation of the data. For example, direct effects of AOM on BBB hyperpermeability have been suggested using a monolayer of mouse brain microvascular endothelia cells¹⁴¹, which has not been confirmed in other studies^{37,142}. However, *in vivo*, the hyperpermeability of the BBB is either not observed¹⁴³ or if it is, it is not until the later stages of HE^{37,142-144} and requires the presence of systemic proinflammatory signals^{37,145}, suggesting that direct effects of AOM on the permeability of the BBB is negligible *in vivo*. APAP has been demonstrated to have a direct toxic effect on the brain in rats^{22,25} and issues may be present in other toxin models as well and therefore experiments should be designed to consider these possible off target effects. Secondly, issues with reproducibility of these models, both between laboratories and between experiments within a single laboratory are often observed. These variations may arise due to a multitude of reasons including animal strains. It has been clearly demonstrated that BDL Wistar rats can be studied at longer time points (8 weeks) than BDL Sprague-Dawley rats (6 weeks)^{104,105,109}. Other factors include variations in lot quality of the toxin, environmental differences, male vs female, as well as alterations in standard animal husbandry and nature of the supportive care provided. Variations between

Accepted Article
experiments within one lab may be somewhat mitigated by careful experimental design, and strict adherence to the supportive care regime (e.g monitoring body temperature, dextrose supplementation to prevent hypoglycemia etc¹⁴³). Differences in outcomes between laboratories are a little more difficult to control for, although a thorough description of the experimental conditions, to include strain details, analgesics, and other supportive care strategies used should be included in publications to aide in the reproducibility of the experimental model.

Future Directions:

Even though a substantial amount of new evidence and insights have emerged from using the above-mentioned animal models of HE, some aspects still remain to be developed or improved on. Specifically;

1. There remains a distinct lack of CLD models that progress to HE and mirror the common etiologies of HE in humans (e.g., alcoholic liver disease, viral hepatitis and NAFLD). These etiological factors may exhibit features of mild fibrosis but fail to exhibit features of severe liver fibrosis or cirrhosis in animal models. The impact of etiological factors on HE merits to be investigated.
2. Many existing models of liver fibrosis, in particular genetic models such as the MDR2 knockout mouse or the PDGF-B transgenic mouse, have not yet been fully characterized for features of HE¹⁴⁶⁻¹⁴⁸.
3. Cholestatic liver disease affects both children and adults. In children, there is emerging evidence that chronic cholestasis early in life may be associated with long-term neurocognitive and neuromotor impairment¹⁴⁹⁻¹⁵¹, but studies examining the effect of liver disease during brain development are missing. In this context, ligation of the bile-duct can also be performed in young rats (i.e. 15 or 21 days after birth) leading to a model of type C HE in the developing brain¹⁵².
4. Differences between sex in children with biliary atresia have been observed¹⁵³, with more women presenting with cholestatic liver disease than men. In this context, sex differences in animal models of HE warrants further investigation.

5. Cognitive tests including those with increased sensitivity (touchscreen cognitive testing) merit to be implicated in the evaluation of HE in rodents (Figure 3). Electroencephalography (EEG) in rodents also warrants further studies¹⁵⁴.
6. Monitoring the brain using magnetic resonance imaging, magnetic resonance spectroscopy, microdialysis, positron emission tomography are excellent tools to elucidate the relationship between liver function, brain metabolic alterations, cellular changes, cell swelling/edema, and neurological manifestations in HE. Further longitudinal, multiparametric and multimodal studies are warranted.

Recommendations of the ISHEN working group

Numerous different animal models of HE currently exist and have been well-characterized. Based on practicality and reproducibility, certain models are recommended (Figure 4). These recommendations were based on the consensus agreement of our working group and are based on ease and reproducibility of the liver disease, the development of HE-associated neurological deficits and hyperammonemia. We have not based these recommendations on other features that may be associated with HE pathogenesis (e.g. biochemical analyses, or imaging studies) as these have not been consistently studied in the majority of models. However, an animal model that exactly mirrors the etiology and subsequent effects on the brain in humans does not exist, highlighting the need for further refinement of existing models and the development of novel animal models for HE. New models should be developed with a particular emphasis on both the etiology (for example a rodent model of end-stage alcoholic liver disease resulting in HE), the underlying liver pathology and the pathogenic features that are associated with the development of each type of HE. To this end, key features that should be characterized when developing and classifying a novel animal model of HE include: i) type of underlying liver pathology, ii) hyperammonemia and iii) cognitive and neurological deficits.

In conclusion, limitations are found of each of the current HE models just as no animal model exists that replicates the full human condition. However, animal models are critical for

answering specific pathophysiological questions. They are also valuable in allowing to initially investigate, understand and test novel therapeutic strategies which could not be conducted in humans. It is therefore vital to choose the appropriate model when considering key features of HE to be studied.

Legends:

Table 1: Neurological score assessment in acute liver failure in mice

A categorical neurological scoring system based on 5 individual reflexes and presence of ataxia in mice. Neurological score is determined by assigning a semi-quantitative evaluation to each of the above-mentioned parameters. 0 (no reflex evident/significant ataxia), 1 (weak or delayed reflex/minor ataxia), or 2 (intact reflex/no ataxia). The summation of these six reflexes and ataxia gives a neurological score between 0 and 12. This scoring system has been developed for the assessment of acute liver failure in mice. The applicability of this scale to rat models of acute liver failure are lacking

Table 2: Examples of behavioral tests and main findings in different models of type B and C HE.

Figure 1: Common methods for the induction of liver failure/disease

Various ways of inducing acute liver failure or chronic liver disease. Red boxes; surgical methods, grey boxes; toxin approaches.

Figure 2: Advantages and disadvantages of working with different experimental animals.

Figure 3: Behavioral tests for investigating HE in rodents

Various and numerous tests or methods are available to evaluate cognitive impairment, sensory-motor deficits as well as anxiety and depression.

Figure 4: Recommendations for Type A, B and C HE animal models

The practicality, reproducibility and recommendations for the different animal models of type A, B and C HE. *** Excellent, ** Moderate * Poor.

References

1. Cabrera-Pastor A, Llansola M, Montoliu C, et al. Peripheral inflammation induces neuroinflammation that alters neurotransmission and cognitive and motor function in hepatic encephalopathy: Underlying mechanisms and therapeutic implications. *Acta Physiol (Oxf)*. 2019;226(2):e13270.
2. Bosoi CR, Rose CF. Oxidative stress: a systemic factor implicated in the pathogenesis of hepatic encephalopathy. *Metab Brain Dis*. 2013;28(2):175-178.
3. McMillin M, Frampton G, Thompson M, et al. Neuronal CCL2 is upregulated during hepatic encephalopathy and contributes to microglia activation and neurological decline. *J Neuroinflammation*. 2014;11:121.
4. McMillin M, Grant S, Frampton G, Andry S, Brown A, DeMorrow S. Fractalkine suppression during hepatic encephalopathy promotes neuroinflammation in mice. *J Neuroinflammation*. 2016;13(1):198.
5. McMillin M, Grant S, Frampton G, et al. Elevated circulating TGFbeta1 during acute liver failure activates TGFbetaR2 on cortical neurons and exacerbates neuroinflammation and hepatic encephalopathy in mice. *J Neuroinflammation*. 2019;16(1):69.
6. Matkowskyj KA, Marrero JA, Carroll RE, Danilkovich AV, Green RM, Benya RV. Azoxymethane-induced fulminant hepatic failure in C57BL/6J mice: characterization of a new animal model. *Am J Physiol*. 1999;277(2):G455-462.
7. McMillin M, Galindo C, Pae HY, et al. Gli1 activation and protection against hepatic encephalopathy is suppressed by circulating transforming growth factor beta1 in mice. *J Hepatol*. 2014;61(6):1260-1266.
8. Bemeur C, Cudalbu C, Dam G, Thrane AS, Cooper AJ, Rose CF. Brain edema: a valid endpoint for measuring hepatic encephalopathy? *Metab Brain Dis*. 2016;31(6):1249-1258.
9. Bosoi CR, Rose CF. Brain edema in acute liver failure and chronic liver disease: similarities and differences. *Neurochem Int*. 2013;62(4):446-457.

- Accepted Article
10. Bajaj JS, Bloom PP, Chung RT, et al. Variability and Lability of Ammonia Levels in Healthy Volunteers and Patients With Cirrhosis: Implications for Trial Design and Clinical Practice. *Am J Gastroenterol*. 2020;115(5):783-785.
 11. Hashim IA, Cuthbert JA. Elevated ammonia concentrations: potential for pre-analytical and analytical contributing factors. *Clin Biochem*. 2014;47(16-17):233-236.
 12. El-Khoury JM, Bunch DR, Wang S. Is the effect of hemolysis on plasma ammonia measurement overrated? *Arch Pathol Lab Med*. 2012;136(5):471-472.
 13. Goldstein BN, Wesler J, Nowacki AS, Reineks E, Natowicz MR. Investigations of blood ammonia analysis: Test matrices, storage, and stability. *Clin Biochem*. 2017;50(9):537-539.
 14. de Graaf RA, Chowdhury GM, Brown PB, Rothman DL, Behar KL. In situ 3D magnetic resonance metabolic imaging of microwave-irradiated rodent brain: a new tool for metabolomics research. *J Neurochem*. 2009;109(2):494-501.
 15. De Chiara F, Thomsen KL, Habtesion A, et al. Ammonia Scavenging Prevents Progression of Fibrosis in Experimental Nonalcoholic Fatty Liver Disease. *Hepatology*. 2020;71(3):874-892.
 16. Ytrebo LM, Kristiansen RG, Maehre H, et al. L-ornithine phenylacetate attenuates increased arterial and extracellular brain ammonia and prevents intracranial hypertension in pigs with acute liver failure. *Hepatology*. 2009;50(1):165-174.
 17. Butterworth RF, Norenberg MD, Felipe V, et al. Experimental models of hepatic encephalopathy: ISHEN guidelines. *Liver Int*. 2009;29(6):783-788.
 18. McGill MR, Williams CD, Xie Y, Ramachandran A, Jaeschke H. Acetaminophen-induced liver injury in rats and mice: comparison of protein adducts, mitochondrial dysfunction, and oxidative stress in the mechanism of toxicity. *Toxicol Appl Pharmacol*. 2012;264(3):387-394.
 19. Shah N, Montes de Oca M, Jover-Cobos M, et al. Role of toll-like receptor 4 in mediating multiorgan dysfunction in mice with acetaminophen induced acute liver failure. *Liver Transpl*. 2013;19(7):751-761.

20. Panatto JP, Jeremias IC, Ferreira GK, et al. Inhibition of mitochondrial respiratory chain in the brain of rats after hepatic failure induced by acetaminophen. *Mol Cell Biochem.* 2011;350(1-2):149-154.
21. Saad MA, Rastanawi AA, El-Yamany MF. Alogliptin abates memory injuries of hepatic encephalopathy induced by acute paracetamol intoxication via switching-off autophagy-related apoptosis. *Life Sci.* 2018;215:11-21.
22. Vigo MB, Perez MJ, De Fino F, et al. Acute acetaminophen intoxication induces direct neurotoxicity in rats manifested as astrogliosis and decreased dopaminergic markers in brain areas associated with locomotor regulation. *Biochem Pharmacol.* 2019;170:113662.
23. Ilic S, Drmic D, Zarkovic K, et al. High hepatotoxic dose of paracetamol produces generalized convulsions and brain damage in rats. A counteraction with the stable gastric pentadecapeptide BPC 157 (PL 14736). *J Physiol Pharmacol.* 2010;61(2):241-250.
24. Isobe-Harima Y, Terai S, Miura I, et al. A new hepatic encephalopathy model to monitor the change of neural amino acids and astrocytes with behaviour disorder. *Liver Int.* 2008;28(1):117-125.
25. Posadas I, Santos P, Blanco A, Munoz-Fernandez M, Cena V. Acetaminophen induces apoptosis in rat cortical neurons. *PLoS One.* 2010;5(12):e15360.
26. Heidari R, Jamshidzadeh A, Niknahad H, et al. Effect of taurine on chronic and acute liver injury: Focus on blood and brain ammonia. *Toxicol Rep.* 2016;3:870-879.
27. Scorticati C, Prestifilippo JP, Eizayaga FX, et al. Hyperammonemia, brain edema and blood-brain barrier alterations in prehepatic portal hypertensive rats and paracetamol intoxication. *World J Gastroenterol.* 2004;10(9):1321-1324.
28. Newsome PN, Plevris JN, Nelson LJ, Hayes PC. Animal models of fulminant hepatic failure: a critical evaluation. *Liver Transpl.* 2000;6(1):21-31.
29. Thiel C, Thiel K, Etspueler A, et al. A reproducible porcine model of acute liver failure induced by intrajejunal acetaminophen administration. *Eur Surg Res.* 2011;46(3):118-126.

- Accepted Article
30. Lee KC, Palacios Jimenez C, Alibhai H, et al. A reproducible, clinically relevant, intensively managed, pig model of acute liver failure for testing of therapies aimed to prolong survival. *Liver Int.* 2013;33(4):544-551.
 31. Scheuermann K, Thiel C, Thiel K, et al. Correlation of the intracranial pressure to the central venous pressure in the late phase of acute liver failure in a porcine model. *Acta Neurochir Suppl.* 2012;114:387-391.
 32. Baker LA, Lee KC, Palacios Jimenez C, et al. Circulating microRNAs Reveal Time Course of Organ Injury in a Porcine Model of Acetaminophen-Induced Acute Liver Failure. *PLoS One.* 2015;10(5):e0128076.
 33. Lee KCL, Baker L, Mallett S, et al. Hypercoagulability progresses to hypocoagulability during evolution of acetaminophen-induced acute liver injury in pigs. *Sci Rep.* 2017;7(1):9347.
 34. Belanger M, Cote J, Butterworth RF. Neurobiological characterization of an azoxymethane mouse model of acute liver failure. *Neurochem Int.* 2006;48(6-7):434-440.
 35. Hori T, Chen F, Baine AM, Gardner LB, Nguyen JH. Fulminant liver failure model with hepatic encephalopathy in the mouse. *Ann Gastroenterol.* 2011;24(4):294-306.
 36. Megaraj V, Ding X, Fang C, Kovalchuk N, Zhu Y, Zhang QY. Role of hepatic and intestinal p450 enzymes in the metabolic activation of the colon carcinogen azoxymethane in mice. *Chem Res Toxicol.* 2014;27(4):656-662.
 37. McMillin MA, Frampton GA, Seiwel AP, Patel NS, Jacobs AN, DeMorrow S. TGFbeta1 exacerbates blood-brain barrier permeability in a mouse model of hepatic encephalopathy via upregulation of MMP9 and downregulation of claudin-5. *Lab Invest.* 2015;95(8):903-913.
 38. Chirito E, Reiter B, Lister C, Chang TM. Artificial liver: the effect of ACAC microencapsulated charcoal hemoperfusion on fulminant hepatic failure. *Artif Organs.* 1977;1(1):76-83.

39. Dixit V, Chang TM. Brain edema and the blood brain barrier in galactosamine-induced fulminant hepatic failure rats. An animal model for evaluation of liver support systems. *ASAIO Trans.* 1990;36(1):21-27.
40. Keppler D, Lesch R, Reutter W, Decker K. Experimental hepatitis induced by D-galactosamine. *Exp Mol Pathol.* 1968;9(2):279-290.
41. Cauli O, Gonzalez-Usano A, Cabrera-Pastor A, et al. Blocking NMDA receptors delays death in rats with acute liver failure by dual protective mechanisms in kidney and brain. *Neuromolecular Med.* 2014;16(2):360-375.
42. Glorioso JM, Mao SA, Rodysill B, et al. Pivotal preclinical trial of the spheroid reservoir bioartificial liver. *J Hepatol.* 2015;63(2):388-398.
43. van de Kerkhove MP, Hoekstra R, van Gulik TM, Chamuleau RA. Large animal models of fulminant hepatic failure in artificial and bioartificial liver support research. *Biomaterials.* 2004;25(9):1613-1625.
44. Muller A, Machnik F, Zimmermann T, Schubert H. Thioacetamide-induced cirrhosis-like liver lesions in rats--usefulness and reliability of this animal model. *Exp Pathol.* 1988;34(4):229-236.
45. Hilgier W, Olson JE. Brain ion and amino acid contents during edema development in hepatic encephalopathy. *J Neurochem.* 1994;62(1):197-204.
46. Rama Rao KV, Reddy PV, Tong X, Norenberg MD. Brain edema in acute liver failure: inhibition by L-histidine. *Am J Pathol.* 2010;176(3):1400-1408.
47. Zimmermann C, Ferenci P, Pifl C, et al. Hepatic encephalopathy in thioacetamide-induced acute liver failure in rats: characterization of an improved model and study of amino acid-ergic neurotransmission. *Hepatology.* 1989;9(4):594-601.
48. Abdel-Rafei M, Amin MM, Hasan HF. Novel effect of Daflon and low-dose gamma-radiation in modulation of thioacetamide-induced hepatic encephalopathy in male albino rats. *Hum Exp Toxicol.* 2017;36(1):62-81.
49. Jayakumar AR, Valdes V, Norenberg MD. The Na-K-Cl cotransporter in the brain edema of acute liver failure. *J Hepatol.* 2011;54(2):272-278.

50. Tsai CY, Su CH, Chan JY, Chan SH. Nitrosative Stress-Induced Disruption of Baroreflex Neural Circuits in a Rat Model of Hepatic Encephalopathy: A DTI Study. *Sci Rep*. 2017;7:40111.
51. Sathyasaikumar KV, Swapna I, Reddy PV, et al. Co-administration of C-Phycocyanin ameliorates thioacetamide-induced hepatic encephalopathy in Wistar rats. *J Neurol Sci*. 2007;252(1):67-75.
52. Jia W, Liu J, Hu R, et al. Xiaochaihutang Improves the Cortical Astrocyte Edema in Thioacetamide-Induced Rat Acute Hepatic Encephalopathy by Activating NRF2 Pathway. *Front Pharmacol*. 2020;11:382.
53. Itzhak Y, Roig-Cantisano A, Dombro RS, Norenberg MD. Acute liver failure and hyperammonemia increase peripheral-type benzodiazepine receptor binding and pregnenolone synthesis in mouse brain. *Brain Res*. 1995;705(1-2):345-348.
54. Jayakumar AR, Bethea JR, Tong XY, Gomez J, Norenberg MD. NF-kappaB in the mechanism of brain edema in acute liver failure: studies in transgenic mice. *Neurobiol Dis*. 2011;41(2):498-507.
55. Sarhan S, Knodgen B, Grauffel C, Seiler N. Effects of inhibition of ornithine aminotransferase on thioacetamide-induced hepatogenic encephalopathy. *Neurochem Res*. 1993;18(4):539-549.
56. Bai Y, Wang S, Wu F, Xie X, Wang Y, Yang Y. The Changes of Mitochondria in Substantia Nigra and Anterior Cerebral Cortex of Hepatic Encephalopathy Induced by Thioacetamide. *Anat Rec (Hoboken)*. 2019;302(7):1169-1177.
57. Chung C, Gottstein J, Blei AT. Indomethacin prevents the development of experimental ammonia-induced brain edema in rats after portacaval anastomosis. *Hepatology*. 2001;34(2):249-254.
58. Swain M, Butterworth RF, Blei AT. Ammonia and related amino acids in the pathogenesis of brain edema in acute ischemic liver failure in rats. *Hepatology*. 1992;15(3):449-453.

59. Traber P, DalCanto M, Ganger D, Blei AT. Effect of body temperature on brain edema and encephalopathy in the rat after hepatic devascularization. *Gastroenterology*. 1989;96(3):885-891.
60. Chatauret N, Rose C, Therrien G, Butterworth RF. Mild hypothermia prevents cerebral edema and CSF lactate accumulation in acute liver failure. *Metab Brain Dis*. 2001;16(1-2):95-102.
61. Rose C, Michalak A, Pannunzio M, Chatauret N, Rambaldi A, Butterworth RF. Mild hypothermia delays the onset of coma and prevents brain edema and extracellular brain glutamate accumulation in rats with acute liver failure. *Hepatology*. 2000;31(4):872-877.
62. Jiang W, Desjardins P, Butterworth RF. Direct evidence for central proinflammatory mechanisms in rats with experimental acute liver failure: protective effect of hypothermia. *J Cereb Blood Flow Metab*. 2009;29(5):944-952.
63. Chastre A, Jiang W, Desjardins P, Butterworth RF. Ammonia and proinflammatory cytokines modify expression of genes coding for astrocytic proteins implicated in brain edema in acute liver failure. *Metab Brain Dis*. 2010;25(1):17-21.
64. Ryska M, Kieslichova E, Pantoflicek T, et al. Devascularization surgical model of acute liver failure in minipigs. *Eur Surg Res*. 2004;36(3):179-184.
65. Ytrebo LM, Ekse S, Sen S, et al. Contractile response of femoral arteries in pigs with acute liver failure. *Scand J Gastroenterol*. 2004;39(10):1000-1004.
66. Wang H, Ohkohchi N, Enomoto Y, et al. Effect of portocaval shunt on residual extreme small liver after extended hepatectomy in porcine. *World J Surg*. 2006;30(11):2014-2022; discussion 2023-2014.
67. Arkadopoulos N, Defterevos G, Nastos C, et al. Development of a porcine model of post-hepatectomy liver failure. *J Surg Res*. 2011;170(2):e233-242.
68. Ladurner R, Schenk M, Margreiter R, Offner F, Konigsrainer A. Influence of portosystemic shunt on liver regeneration after hepatic resection in pigs. *HPB Surg*. 2009;2009:835965.

69. Ladurner R, Traub F, Schenk M, Konigsrainer A, Glatzle J. Cellular liver regeneration after extended hepatic resection in pigs. *HPB Surg.* 2009;2009:306740.
70. Molino G, Avagnina P, Garrone C, et al. Time-dependent modifications of splanchnic circulation in female pigs submitted to end-to-side portacaval anastomosis. *Dig Dis Sci.* 2001;46(3):489-494.
71. Ytrebo LM, Nedredal GI, Langbakk B, Revhaug A. An experimental large animal model for the assessment of bioartificial liver support systems in fulminant hepatic failure. *Scand J Gastroenterol.* 2002;37(9):1077-1088.
72. Kristiansen RG, Lindal S, Myreng K, Revhaug A, Ytrebo LM, Rose CF. Neuropathological changes in the brain of pigs with acute liver failure. *Scand J Gastroenterol.* 2010;45(7-8):935-943.
73. Rose C, Ytrebo LM, Davies NA, et al. Association of reduced extracellular brain ammonia, lactate, and intracranial pressure in pigs with acute liver failure. *Hepatology.* 2007;46(6):1883-1892.
74. Sen S, Rose C, Ytrebo LM, et al. Effect of albumin dialysis on intracranial pressure increase in pigs with acute liver failure: a randomized study. *Crit Care Med.* 2006;34(1):158-164.
75. Nedredal GI, Elvevold K, Ytrebo LM, et al. Significant contribution of liver nonparenchymal cells to metabolism of ammonia and lactate and cocultivation augments the functions of a bioartificial liver. *Am J Physiol Gastrointest Liver Physiol.* 2007;293(1):G75-83.
76. Yagi S, Iida T, Hori T, et al. Effect of portal haemodynamics on liver graft and intestinal mucosa after small-for-size liver transplantation in swine. *Eur Surg Res.* 2012;48(3):163-170.
77. Cudalbu C, McLin VA, Lei H, et al. The C57BL/6J mouse exhibits sporadic congenital portosystemic shunts. *PLoS One.* 2013;8(7):e69782.
78. Or M, Devriendt N, Kitshoff AM, et al. Ammonia concentrations in arterial blood, venous blood, and cerebrospinal fluid of dogs with and without congenital extrahepatic portosystemic shunts. *Am J Vet Res.* 2017;78(11):1313-1318.

79. Bosoi CR, Tremblay M, Rose CF. Induction of systemic oxidative stress leads to brain oedema in portacaval shunted rats. *Liver Int.* 2014;34(9):1322-1329.
80. Gandhi CR, Murase N, Subbotin VM, et al. Portacaval shunt causes apoptosis and liver atrophy in rats despite increases in endogenous levels of major hepatic growth factors. *J Hepatol.* 2002;37(3):340-348.
81. Apelqvist G, Hindfelt B, Andersson G, Bengtsson F. Altered adaptive behaviour expressed in an open-field paradigm in experimental hepatic encephalopathy. *Behav Brain Res.* 1999;106(1-2):165-173.
82. Coy DL, Mehta R, Zee P, Salchli F, Turek FW, Blei AT. Portal-systemic shunting and the disruption of circadian locomotor activity in the rat. *Gastroenterology.* 1992;103(1):222-228.
83. Cuschieri A, Baker PR, Holley MP, Hanson C. Portacaval shunt in the pig 1. Effect on survival, behavior, nutrition, and liver function. *J Surg Res.* 1974;17(6):387-396.
84. Halvorsen JF, Myking AO. Prehepatic portal hypertension in the rat. Immediate and long-term effects on portal vein and aortic pressure of a graded portal vein stenosis, followed by occlusion of the portal vein and spleno-renal collaterals. *Eur Surg Res.* 1979;11(2):89-98.
85. Praktijnjo M, Simon-Talero M, Romer J, et al. Total area of spontaneous portosystemic shunts independently predicts hepatic encephalopathy and mortality in liver cirrhosis. *J Hepatol.* 2020.
86. Rokusz A, Veres D, Szucs A, et al. Ductular reaction correlates with fibrogenesis but does not contribute to liver regeneration in experimental fibrosis models. *PLoS One.* 2017;12(4):e0176518.
87. Yang N, Liu H, Jiang Y, et al. Lactulose enhances neuroplasticity to improve cognitive function in early hepatic encephalopathy. *Neural Regen Res.* 2015;10(9):1457-1462.
88. Liu R, Kang JD, Sartor RB, et al. Neuroinflammation in Murine Cirrhosis Is Dependent on the Gut Microbiome and Is Attenuated by Fecal Transplant. *Hepatology.* 2019.
89. Kang DJ, Betrapally NS, Ghosh SA, et al. Gut microbiota drive the development of neuroinflammatory response in cirrhosis in mice. *Hepatology.* 2016;64(4):1232-1248.

90. Liu R, Ahluwalia V, Kang JD, et al. Effect of Increasing Age on Brain Dysfunction in Cirrhosis. *Hepatol Commun*. 2019;3(1):63-73.
91. Vairappan B, Sundhar M, Srinivas BH. Resveratrol Restores Neuronal Tight Junction Proteins Through Correction of Ammonia and Inflammation in CCl4-Induced Cirrhotic Mice. *Mol Neurobiol*. 2019;56(7):4718-4729.
92. Nicaise C, Prozzi D, Viaene E, et al. Control of acute, chronic, and constitutive hyperammonemia by wild-type and genetically engineered *Lactobacillus plantarum* in rodents. *Hepatology*. 2008;48(4):1184-1192.
93. Haeger P, Bouchet A, Ossandon C, Bresky G. Treatment with Melatonin Improves Cognitive Behavior and Motor Skills in a Rat Model of Liver Fibrosis. *Ann Hepatol*. 2019;18(1):101-108.
94. Schmandra TC, Bauer H, Petrowsky H, Herrmann G, Encke A, Hanisch E. Effect of fibrin glue occlusion of the hepatobiliary tract on thioacetamide-induced liver failure. *Am J Surg*. 2001;182(1):58-63.
95. Singh S, Trigun SK. Activation of neuronal nitric oxide synthase in cerebellum of chronic hepatic encephalopathy rats is associated with up-regulation of NADPH-producing pathway. *Cerebellum*. 2010;9(3):384-397.
96. Jayakumar AR, Tong XY, Curtis KM, et al. Decreased astrocytic thrombospondin-1 secretion after chronic ammonia treatment reduces the level of synaptic proteins: in vitro and in vivo studies. *J Neurochem*. 2014;131(3):333-347.
97. Masumi S, Moriyama M, Kannan Y, et al. Characteristics of nitrogen metabolism in rats with thioacetamide-induced liver cirrhosis. *Toxicology*. 1999;132(2-3):155-166.
98. Noda S, Masumi S, Moriyama M, et al. Population of hepatic macrophages and response of perfused liver to platelet-activating factor during production of thioacetamide-induced cirrhosis in rats. *Hepatology*. 1996;24(2):412-418.
99. Wallace MC, Hamesch K, Lunova M, et al. Standard operating procedures in experimental liver research: thioacetamide model in mice and rats. *Lab Anim*. 2015;49(1 Suppl):21-29.

- Accepted Article
100. Kang JS, Wanibuchi H, Morimura K, et al. Enhancement by estradiol 3-benzoate in thioacetamide-induced liver cirrhosis of rats. *Toxicol Sci.* 2005;85(1):720-726.
 101. Li X, Benjamin IS, Alexander B. Reproducible production of thioacetamide-induced macronodular cirrhosis in the rat with no mortality. *J Hepatol.* 2002;36(4):488-493.
 102. Low TY, Leow CK, Salto-Tellez M, Chung MC. A proteomic analysis of thioacetamide-induced hepatotoxicity and cirrhosis in rat livers. *Proteomics.* 2004;4(12):3960-3974.
 103. Balasubramanian V, Mehta G, Jones H, et al. Post-Transcriptional Regulation of Hepatic DDAH1 with TNF Blockade Leads to Improved eNOS Function and Reduced Portal Pressure In Cirrhotic Rats. *Sci Rep.* 2017;7(1):17900.
 104. Bosoi CR, Parent-Robitaille C, Anderson K, Tremblay M, Rose CF. AST-120 (spherical carbon adsorbent) lowers ammonia levels and attenuates brain edema in bile duct-ligated rats. *Hepatology.* 2011;53(6):1995-2002.
 105. Braissant O, Rackayova V, Pierzchala K, Grosse J, McLin VA, Cudalbu C. Longitudinal neurometabolic changes in the hippocampus of a rat model of chronic hepatic encephalopathy. *J Hepatol.* 2019;71(3):505-515.
 106. Leke R, Oliveira DL, Forgiarini LF, et al. Impairment of short term memory in rats with hepatic encephalopathy due to bile duct ligation. *Metab Brain Dis.* 2013;28(2):187-192.
 107. Bosoi CR, Zwingmann C, Marin H, et al. Increased brain lactate is central to the development of brain edema in rats with chronic liver disease. *J Hepatol.* 2014;60(3):554-560.
 108. Leke R, de Oliveira DL, Mussulini BH, et al. Impairment of the organization of locomotor and exploratory behaviors in bile duct-ligated rats. *PLoS One.* 2012;7(5):e36322.
 109. Rackayova V, Braissant O, McLin VA, Berset C, Lanz B, Cudalbu C. (1)H and (31)P magnetic resonance spectroscopy in a rat model of chronic hepatic encephalopathy: in vivo longitudinal measurements of brain energy metabolism. *Metab Brain Dis.* 2016;31(6):1303-1314.

110. Chavarria L, Oria M, Romero-Gimenez J, Alonso J, Lope-Piedrafita S, Cordoba J. Brain magnetic resonance in experimental acute-on-chronic liver failure. *Liver Int.* 2013;33(2):294-300.
111. Davies NA, Wright G, Ytrebo LM, et al. L-ornithine and phenylacetate synergistically produce sustained reduction in ammonia and brain water in cirrhotic rats. *Hepatology.* 2009;50(1):155-164.
112. Cudalbu C, Taylor-Robinson SD. Brain Edema in Chronic Hepatic Encephalopathy. *J Clin Exp Hepatol.* 2019;9(3):362-382.
113. Rodrigo R, Cauli O, Gomez-Pinedo U, et al. Hyperammonemia induces neuroinflammation that contributes to cognitive impairment in rats with hepatic encephalopathy. *Gastroenterology.* 2010;139(2):675-684.
114. Wright G, Davies NA, Shawcross DL, et al. Endotoxemia produces coma and brain swelling in bile duct ligated rats. *Hepatology.* 2007;45(6):1517-1526.
115. Wright G, Soper R, Brooks HF, et al. Role of aquaporin-4 in the development of brain oedema in liver failure. *J Hepatol.* 2010;53(1):91-97.
116. D'Mello C, Almishri W, Liu H, Swain MG. Interactions Between Platelets and Inflammatory Monocytes Affect Sickness Behavior in Mice With Liver Inflammation. *Gastroenterology.* 2017;153(5):1416-1428 e1412.
117. Nguyen K, D'Mello C, Le T, Urbanski S, Swain MG. Regulatory T cells suppress sickness behaviour development without altering liver injury in cholestatic mice. *J Hepatol.* 2012;56(3):626-631.
118. O'Brien A, China L, Massey KA, et al. Bile duct-ligated mice exhibit multiple phenotypic similarities to acute decompensation patients despite histological differences. *Liver Int.* 2016;36(6):837-846.
119. Therrien G, Rose C, Butterworth J, Butterworth RF. Protective effect of L-carnitine in ammonia-precipitated encephalopathy in the portacaval shunted rat. *Hepatology.* 1997;25(3):551-556.
120. Katz S, Jimenez MA, Lehmkuhler WE, Grosfeld JL. Liver bacterial clearance following hepatic artery ligation and portacaval shunt. *J Surg Res.* 1991;51(3):267-270.

121. Garcia-Lezana T, Oria M, Romero-Gimenez J, et al. Cerebellar neurodegeneration in a new rat model of episodic hepatic encephalopathy. *J Cereb Blood Flow Metab.* 2017;37(3):927-937.
122. Oria M, Chatauret N, Chavarria L, et al. Motor-evoked potentials in awake rats are a valid method of assessing hepatic encephalopathy and of studying its pathogenesis. *Hepatology.* 2010;52(6):2077-2085.
123. Oria M, Romero-Gimenez J, Arranz JA, Riudor E, Raguer N, Cordoba J. Ornithine phenylacetate prevents disturbances of motor-evoked potentials induced by intestinal blood in rats with portacaval anastomosis. *J Hepatol.* 2012;56(1):109-114.
124. Davuluri G, Allawy A, Thapaliya S, et al. Hyperammonaemia-induced skeletal muscle mitochondrial dysfunction results in cataplerosis and oxidative stress. *J Physiol.* 2016;594(24):7341-7360.
125. Davuluri G, Krokowski D, Guan BJ, et al. Metabolic adaptation of skeletal muscle to hyperammonemia drives the beneficial effects of l-leucine in cirrhosis. *J Hepatol.* 2016;65(5):929-937.
126. Harry D, Anand R, Holt S, et al. Increased sensitivity to endotoxemia in the bile duct-ligated cirrhotic Rat. *Hepatology.* 1999;30(5):1198-1205.
127. Azorin I, Minana MD, Felipo V, Grisolia S. A simple animal model of hyperammonemia. *Hepatology.* 1989;10(3):311-314.
128. Wright G, Vairappan B, Stadlbauer V, Mookerjee RP, Davies NA, Jalan R. Reduction in hyperammonaemia by ornithine phenylacetate prevents lipopolysaccharide-induced brain edema and coma in cirrhotic rats. *Liver Int.* 2012;32(3):410-419.
129. Agostoni V, Lee SH, Forster V, et al. Liposome-Supported Peritoneal Dialysis for the Treatment of Hyperammonemia-Associated Encephalopathy. *Adv Funct Mater* 2016;26(46):8382–8389.
130. Rackayova V, Flatt E, Braissant O, et al. Probiotics improve the neurometabolic profile of rats with chronic cholestatic liver disease. *Sci Rep.* 2021;11(1):2269.
131. Johansson M, Agusti A, Llansola M, et al. GR3027 antagonizes GABAA receptor-potentiating neurosteroids and restores spatial learning and motor coordination in rats

- with chronic hyperammonemia and hepatic encephalopathy. *Am J Physiol Gastrointest Liver Physiol*. 2015;309(5):G400-409.
132. Bemeur C, Vaquero J, Desjardins P, Butterworth RF. N-acetylcysteine attenuates cerebral complications of non-acetaminophen-induced acute liver failure in mice: antioxidant and anti-inflammatory mechanisms. *Metab Brain Dis*. 2010;25(2):241-249.
133. Jiang W, Desjardins P, Butterworth RF. Minocycline attenuates oxidative/nitrosative stress and cerebral complications of acute liver failure in rats. *Neurochem Int*. 2009;55(7):601-605.
134. Rose C, Michalak A, Rao KV, Quack G, Kircheis G, Butterworth RF. L-ornithine-L-aspartate lowers plasma and cerebrospinal fluid ammonia and prevents brain edema in rats with acute liver failure. *Hepatology*. 1999;30(3):636-640.
135. Bemeur C, Desjardins P, Butterworth RF. Antioxidant and anti-inflammatory effects of mild hypothermia in the attenuation of liver injury due to azoxymethane toxicity in the mouse. *Metab Brain Dis*. 2010;25(1):23-29.
136. Bemeur C, Qu H, Desjardins P, Butterworth RF. IL-1 or TNF receptor gene deletion delays onset of encephalopathy and attenuates brain edema in experimental acute liver failure. *Neurochem Int*. 2010;56(2):213-215.
137. Chastre A, Belanger M, Beauchesne E, Nguyen BN, Desjardins P, Butterworth RF. Inflammatory cascades driven by tumor necrosis factor-alpha play a major role in the progression of acute liver failure and its neurological complications. *PLoS One*. 2012;7(11):e49670.
138. McMillin M, Frampton G, Grant S, et al. Bile Acid-Mediated Sphingosine-1-Phosphate Receptor 2 Signaling Promotes Neuroinflammation during Hepatic Encephalopathy in Mice. *Front Cell Neurosci*. 2017;11:191.
139. McMillin M, Frampton G, Quinn M, et al. Bile Acid Signaling Is Involved in the Neurological Decline in a Murine Model of Acute Liver Failure. *Am J Pathol*. 2016;186(2):312-323.

140. McMillin M, Grant S, Frampton G, et al. FXR-Mediated Cortical Cholesterol Accumulation Contributes to the Pathogenesis of Type A Hepatic Encephalopathy. *Cell Mol Gastroenterol Hepatol*. 2018;6(1):47-63.
141. Jayakumar AR, Ruiz-Cordero R, Tong XY, Norenberg MD. Brain edema in acute liver failure: role of neurosteroids. *Arch Biochem Biophys*. 2013;536(2):171-175.
142. Grant S, McMillin M, Frampton G, et al. Direct Comparison of the Thioacetamide and Azoxymethane Models of Type A Hepatic Encephalopathy in Mice. *Gene Expr*. 2018;18(3):171-185.
143. Bemeur C, Chastre A, Desjardins P, Butterworth RF. No changes in expression of tight junction proteins or blood–brain barrier permeability in azoxymethane-induced experimental acute liver failure. *Neurochemistry International*. 2010;56(2):205-207.
144. Nguyen JH, Yamamoto S, Steers J, et al. Matrix metalloproteinase-9 contributes to brain extravasation and edema in fulminant hepatic failure mice. *J Hepatol*. 2006;44(6):1105-1114.
145. Chastre A, Belanger M, Nguyen BN, Butterworth RF. Lipopolysaccharide precipitates hepatic encephalopathy and increases blood-brain barrier permeability in mice with acute liver failure. *Liver Int*. 2014;34(3):353-361.
146. Czochra P, Klopčič B, Meyer E, et al. Liver fibrosis induced by hepatic overexpression of PDGF-B in transgenic mice. *J Hepatol*. 2006;45(3):419-428.
147. Thieringer F, Maass T, Czochra P, et al. Spontaneous hepatic fibrosis in transgenic mice overexpressing PDGF-A. *Gene*. 2008;423(1):23-28.
148. Baghdasaryan A, Fickert P, Fuchsbichler A, et al. Role of hepatic phospholipids in development of liver injury in Mdr2 (Abcb4) knockout mice. *Liver Int*. 2008;28(7):948-958.
149. Caudle SE, Katzenstein JM, Karpen SJ, McLin VA. Language and motor skills are impaired in infants with biliary atresia before transplantation. *J Pediatr*. 2010;156(6):936-940 e931.

150. Ohnemus D, Neighbors K, Rychlik K, et al. Health-Related Quality of Life and Cognitive Functioning in Pediatric Liver Transplant Recipients. *Liver Transpl.* 2020;26(1):45-56.
151. Sorensen LG, Neighbors K, Martz K, et al. Longitudinal study of cognitive and academic outcomes after pediatric liver transplantation. *J Pediatr.* 2014;165(1):65-72 e62.
152. Rackayova V, Braissant O, Rougemont AL, Cudalbu C, McLin VA. Longitudinal osmotic and neurometabolic changes in young rats with chronic cholestatic liver disease. *Sci Rep.* 2020;10(1):7536.
153. Caudle SE, Katzenstein JM, Karpen S, McLin V. Developmental assessment of infants with biliary atresia: differences between boys and girls. *J Pediatr Gastroenterol Nutr.* 2012;55(4):384-389.
154. Marini S, Santangeli O, Saarelainen P, et al. Abnormalities in the Polysomnographic, Adenosine and Metabolic Response to Sleep Deprivation in an Animal Model of Hyperammonemia. *Front Physiol.* 2017;8:636.
155. Jover R, Rodrigo R, Felipe V, et al. Brain edema and inflammatory activation in bile duct ligated rats with diet-induced hyperammonemia: A model of hepatic encephalopathy in cirrhosis. *Hepatology.* 2006;43(6):1257-1266.
156. Bosoi CR, Yang X, Huynh J, et al. Systemic oxidative stress is implicated in the pathogenesis of brain edema in rats with chronic liver failure. *Free Radic Biol Med.* 2012;52(7):1228-1235.
157. Dhanda S, Gupta S, Halder A, Sunkaria A, Sandhir R. Systemic inflammation without gliosis mediates cognitive deficits through impaired BDNF expression in bile duct ligation model of hepatic encephalopathy. *Brain Behav Immun.* 2018;70:214-232.
158. Aguilar MA, Minarro J, Felipe V. Chronic moderate hyperammonemia impairs active and passive avoidance behavior and conditional discrimination learning in rats. *Exp Neurol.* 2000;161(2):704-713.
159. Fabri DR, Hott SC, Reis DG, Biojone C, Correa FM, Resstel LB. The expression of contextual fear conditioning involves activation of a NMDA receptor-nitric oxide-cGMP

- pathway in the dorsal hippocampus of rats. *Eur Neuropsychopharmacol*. 2014;24(10):1676-1686.
160. Ji ES, Kim YM, Shin MS, et al. Treadmill exercise enhances spatial learning ability through suppressing hippocampal apoptosis in Huntington's disease rats. *J Exerc Rehabil*. 2015;11(3):133-139.
161. Shen TC, Albenberg L, Bittinger K, et al. Engineering the gut microbiota to treat hyperammonemia. *J Clin Invest*. 2015;125(7):2841-2850.
162. Aghaei I, Saeedi Saravi SS, Ghotbi Ravandi S, et al. Evaluation of prepulse inhibition and memory impairments at early stage of cirrhosis may be considered as a diagnostic index for minimal hepatic encephalopathy. *Physiol Behav*. 2017;173:87-94.
163. Ali SH, Madhana RM, K VA, et al. Resveratrol ameliorates depressive-like behavior in repeated corticosterone-induced depression in mice. *Steroids*. 2015;101:37-42.
164. Bederson JB, Pitts LH, Tsuji M, Nishimura MC, Davis RL, Bartkowski H. Rat middle cerebral artery occlusion: evaluation of the model and development of a neurologic examination. *Stroke*. 1986;17(3):472-476.
165. Bouet V, Freret T, Toutain J, Divoux D, Boulouard M, Schumann-Bard P. Sensorimotor and cognitive deficits after transient middle cerebral artery occlusion in the mouse. *Exp Neurol*. 2007;203(2):555-567.
166. Farrar AM, Murphy CA, Paterson NE, et al. Cognitive deficits in transgenic and knock-in HTT mice parallel those in Huntington's disease. *J Huntingtons Dis*. 2014;3(2):145-158.
167. Fortin NJ, Wright SP, Eichenbaum H. Recollection-like memory retrieval in rats is dependent on the hippocampus. *Nature*. 2004;431(7005):188-191.
168. Franca MER, Ramos R, Oliveira WH, et al. Tadalafil restores long-term memory and synaptic plasticity in mice with hepatic encephalopathy. *Toxicol Appl Pharmacol*. 2019;379:114673.
169. Glynn D, Skillings EA, Morton AJ. A comparison of discrimination learning in touchscreen and 2-choice swim tank using an allelic series of Huntington's disease mice. *J Neurosci Methods*. 2016;265:56-71.

170. Hua Y, Schallert T, Keep RF, Wu J, Hoff JT, Xi G. Behavioral tests after intracerebral hemorrhage in the rat. *Stroke*. 2002;33(10):2478-2484.
171. Lainiola M, Procaccini C, Linden AM. mGluR3 knockout mice show a working memory defect and an enhanced response to MK-801 in the T- and Y-maze cognitive tests. *Behav Brain Res*. 2014;266:94-103.
172. Lee JH, Tigchelaar S, Liu J, et al. Lack of neuroprotective effects of simvastatin and minocycline in a model of cervical spinal cord injury. *Exp Neurol*. 2010;225(1):219-230.
173. Li Y, Mei LH, Qiang JW, Ji CX, Ju S. Reduction of manganese intake improves neuropsychological manifestations in rats with minimal hepatic encephalopathy. *Neuroscience*. 2017;347:148-155.
174. Luo J, Wang T, Liang S, Hu X, Li W, Jin F. Ingestion of Lactobacillus strain reduces anxiety and improves cognitive function in the hyperammonemia rat. *Sci China Life Sci*. 2014;57(3):327-335.
175. Mi XJ, Chen SW, Wang WJ, et al. Anxiolytic-like effect of paeonol in mice. *Pharmacol Biochem Behav*. 2005;81(3):683-687.
176. Torres-Lista V, Lopez-Pousa S, Gimenez-Llort L. Marble-burying is enhanced in 3xTg-AD mice, can be reversed by risperidone and it is modulable by handling. *Behav Processes*. 2015;116:69-74.
177. Wooden JI, Pido J, Mathews H, Kieltyka R, Montemayor BA, Ward CP. Sleep deprivation impairs recall of social transmission of food preference in rats. *Nat Sci Sleep*. 2014;6:129-135.
178. Wu H, Cottingham C, Chen L, et al. Age-dependent differential regulation of anxiety- and depression-related behaviors by neurabin and spinophilin. *PLoS One*. 2017;12(7):e0180638.
179. Yadav R, Gupta SC, Hillman BG, Bhatt JM, Stairs DJ, Dravid SM. Deletion of glutamate delta-1 receptor in mouse leads to aberrant emotional and social behaviors. *PLoS One*. 2012;7(3):e32969.

180. Cauli O, Llansola M, Agusti A, et al. Cerebral oedema is not responsible for motor or cognitive deficits in rats with hepatic encephalopathy. *Liver Int.* 2014;34(3):379-387.


Table 1:

Parameter	Methodology
Pinna reflex	Touching the auditory meatus with a cotton applicator and observing ear retraction or head movement
Corneal reflex	Gently touching the cornea with a cotton applicator and measuring the blink response
Tail flexion	Tail pinch with forceps and assessing tail flexion
Escape response	Tail pinch by forceps and a subsequent movement of mice away from stimuli
Righting reflex	Placing the mice on their backs and assessing the time for them to right themselves
Ataxia	Placing mice on wire cage lid and observing the number of times a foot falls through the lid

Reference	Subjects	Type of test	Results
Leke, et al. ¹⁰⁸	Adult female Wistar rats @ 6 weeks post-BDL	Open Field Elevated Plus-Maze Foot-Fault	Disturbed spontaneous locomotor and exploratory activities as a consequence of altered spatio-temporal organization of behavior
Leke, et al. ⁹⁶	Adult male Wistar rats @ 6 weeks post-BDL	Object Recognition Task	Impaired STM for recognition memory
Jover, et al., ¹⁵⁵	Adult male Wistar rats @ 2 weeks post-BDL and + 3 weeks HD diet	Rotarod Beam Walking Locomotor Activity	Mild impairment of motor coordination ↓ spontaneous activity
Bosoi, et al., ^{104, 156}	Adult male SD rats @ 6 weeks post-BDL BDL + Allopurinol and BDL + AST-120	Locomotor Activity	↓ total distance traveled and normalized after AST-120 or allopurinol
Braissant, et al. ¹⁰⁵	Adult male Wistar rats @ 4, 6, 8 weeks post-BDL	Open Field (Locomotor Activity)	↓ total distance traveled
Dhanda, et al., ¹⁵⁷	Adult male Wistar rats @ 4 weeks post-BDL	Morris Water Maze Memory Retrieval	↑ in total time taken and distance travelled, no change in velocity ↓ no of entries, time spent and distance travelled in platform zone


		Novel Object Recognition Task	↓ no of entries in known and novel object zone
Rodrigo, et al., ¹¹³	Adult male Wistar rats @ 2 weeks post-BDL + 10 and 21 days after ibuprofen	Motor Activity Y-Maze Learning Test	↓ Motor activity ↓ Learning ability Ibuprofen restored learning and motor activity in BDL rats
Cauli, et al., ¹⁸⁰	Male Wistar rats @ 4 weeks post PCS surgery	Automated actimeter (Motor activity) Rotarod and beam walking test (Motor coordination) Y maze test (Learning) Morris water maze (Spatial memory)	↓ Spontaneous motor activity at 4 weeks post-PCS Motor coordination was impaired at 4 weeks post-PCS ↓ Learning ability at 4 weeks post-PCS ↓ Memory at 4 weeks post-PCS

Fig 1:




liv_14911_f1.jpg

Fig 2:



liv_14911_f2.jpg


Fig 3:



liv_14911_f3.jpg

Fig 4:

		Practicality	Reproducibility	Recommended
Type A				
APA	Mice	***	**	**
	Rats	***	**	*
	Pigs	**	**	***
AOM	Mice	***	***	***
	Rats	***	*	*
D-GAL	Mice	**	*	*
	Rats	**	*	*
TAA	Mice	***	**	**
	Rats	***	**	**
HAL	Rats	*	***	***
	Pigs	*	***	***
Type B				
PCA	Rats	*	***	***
	Pigs	*	***	***
Type C				
CCL4	Mice	***	**	**
	Rats	***	**	*
TAA	Rats	***	*	**
BDL	Mice	**	*	*
	Rats	**	***	***



liv_14911_f4.jpg